



University of Minho
School of Engineering

Synthetic biology strategies for the production of plant polyphenolic compounds

Joana L. Rodrigues*, Daniela Gomes, João Rainha, Lígia R. Rodrigues

Centre of Biological Engineering, University of Minho, Campus de Gualtar,
4710-057 Braga, Portugal
* Corresponding author: joanarodrigues@ceb.uminho.pt



CENTRE OF BIOLOGICAL ENGINEERING

Introduction

Polyphenols biological activities

- Antioxidant
- Anti-inflammatory
- Anticancer
- Would healing
- Skin treatment disorders
- Anti-viral
- Anti-septic
- Anti-aging
- Anti-diabetic
- ...



Extraction from plants


- Low amount
- Low yields and purity
- Limited by seasonality
- Limited by pests and extreme weather
- High land / water investment
- Environmentally unfriendly process
- Expensive downstream process



SOLUTION:
SYNTHETIC BIOLOGY




Polyphenols Market:
\$ 2.26 billion by 2027




Results


Ferulic acid production from tyrosine



Curcumin production from ferulic acid




Curcuminoids production in co-culture




Module 1
Co-culture 1:
pCDFduet_TAL + pRSFduet_C3H_COMT
Co-culture 2:
pRSFduet_TAL_C3H + pACYCduet_COMT


Co-culture population was followed due to curcuminoids production and by blue-white screening due to *lacZ* disruption using **CRISPR-Cas9**



Curcuminoids production in mono-culture



Coumarins production



pAC-4CL + pET28-GST-C2'H (+ pCDFduet_TAL)

Conclusions

Module 1: [Ferulic acid] = 1325 μM
Module 2: [Curcumin] = 1530 μM
Mono-culture: [Curcuminoids] = 19 μM
Co-culture: [Curcuminoids] = 126 μM

Highest productions reported so far

[Umbelliferone] = 263 μM

Acronyms

TAL: tyrosine ammonia lyase from *Rhodotorula glutinis*
C3H: 4-coumarate 3-hydroxylase from *Saccharothrix espanaensis*
COMT: caffeic acid O-methyltransferase from *Arabidopsis thaliana*
4CL: 4-coumarate-CoA ligase from *A. thaliana*
DCS: diketide-CoA synthase from *Curcuma longa*
CURS 1: curcumin synthase 1 from *C. longa*
C2'H: coumaroyl-CoA 2'-hydroxylase from *Ipomoea batatas*
GST: glutathione-S-transferase

References

- [1] ProficientMarket. Global Polyphenols Market Will Register a CAGR of Around 9.0% by 2027. [https://proficientmarket.com/press-release/1225/global-polyphenols-market-\(2020\).](https://proficientmarket.com/press-release/1225/global-polyphenols-market-(2020).)
- [2] Rodrigues JL, Rodrigues LR (2020) Biosynthesis and Heterologous Production of Furanocoumarins: Perspectives and Current Challenges. *Natural Product Reports*, Accepted.
- [3] Rodrigues JL, Prather KLJ, Kluskens LD, Rodrigues LR (2015) Heterologous production of Curcuminoids, *Microbiology and Molecular Biology Reviews*, 79, 39-60.
- [4] Rodrigues JL, Gomes D, Rodrigues LR (2020) A combinatorial approach to optimize the production of curcuminoids from tyrosine in *Escherichia coli*, *Frontiers in Bioengineering and Biotechnology*, 59, 1-15.

This study was supported by FCT under the scope of the strategic funding of UID/BIO/04469/2020 unit and BioTecNorte operation (NORTE-01-0145-FEDER-000004) funded by the European Regional Development Fund under the scope of Norte2020 - Programa Operacional Regional do Norte. JR is recipient of a fellowship supported by a doctoral advanced training (SFRH/BD/138325/2018) funded by FCT.